



Important Safety Information

Prescribing Information

For the initial and long-term treatment of Parkinson's disease: Introducing once-daily MIRAPEX ER.

Once a day and you're on your way.

Once-daily MIRAPEX ER may be right for you.

- Proven effective in relieving symptoms of Parkinson's disease
- MIRAPEX ER is taken just once a day, which many people may find more convenient than medicines taken multiple times a day
- MIRAPEX ER contains the same medicine as Mirapex® (pramipexole dihydrochloride) tablets. MIRAPEX has been a trusted treatment for Parkinson's disease for more than 12 years

Only your doctor can determine if MIRAPEX ER is an appropriate choice for you. Your doctor may make changes to your MIRAPEX ER dose to make sure you are taking the right amount of medicine to help control your symptoms.

Already taking MIRAPEX?

If you are currently taking MIRAPEX, you may be able to switch overnight from MIRAPEX to MIRAPEX ER. Your doctor will work with you to make sure you are taking the right dose to help control your symptoms. Be sure you do not take MIRAPEX and MIRAPEX ER together.

UP TO \$40 SAVINGS PER PRESCRIPTION*

Ready to begin treatment with MIRAPEX ER?

Register today, and if you are eligible, you will save up to \$40 on each of your next 4 prescriptions.*

Talk to your doctor to see if MIRAPEX ER is right for you.

REGISTER NOW

*Up to 4 prescriptions.



If you can't afford your MIRAPEX ER, our patient assistance program may be able to help. Call 1-800-556-8317

Indication:

MIRAPEX ER and MIRAPEX tablets are prescription medicines to treat the signs and symptoms of Parkinson's disease (PD).

Important Safety Information:

MIRAPEX ER or MIRAPEX may cause you to fall asleep without warning during daily activities, including talking, eating, and driving, which may result in accidents. Tell your doctor if you feel new or increased sleepiness while taking MIRAPEX ER or MIRAPEX. Do not drive a car, operate a machine, or do anything that needs you to be alert until you know how MIRAPEX ER or MIRAPEX affects you.

Before taking MIRAPEX ER or MIRAPEX, talk to your doctor about all of your medical conditions, including if you feel sleepy during the day. Tell your doctor if you drink alcohol or take other medications that make you drowsy, as these can increase the chance that MIRAPEX ER or MIRAPEX will make you feel sleepy or fall asleep when you should be awake.

When taking MIRAPEX ER or MIRAPEX, hallucinations (unreal visions, sounds or sensations) may occur and you may sometimes feel dizzy, nauseated, faint or sweaty when you sit up or stand quickly.

Some patients taking certain medicines to treat PD, including MIRAPEX ER and MIRAPEX, have reported problems, such as gambling, compulsive eating, compulsive buying, and increased sex drive. If you or your family members notice that you are developing unusual urges or behaviors, talk to your doctor.

The most common side effects in people taking MIRAPEX ER for early PD are sleepiness, nausea and vomiting, constipation, dizziness, tiredness, hallucinations, dry mouth, muscle spasms, and edema (swelling of the feet and ankles).

The most common side effects in people taking MIRAPEX ER who have later-stage PD are abnormal movements, nausea, constipation, hallucinations, headache, and anorexia.

The most common side effects in people taking MIRAPEX for PD are nausea, dizziness, sleepiness, constipation, hallucinations, insomnia, muscle weakness, confusion, and abnormal movements.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full Prescribing Information for [MIRAPEX ER](#) and [MIRAPEX](#).

This information is intended for U.S. residents only.

Global: added MIRAPEX PI and "US residents" note.



Once daily

global change to "for initial and long-term ..." instead of "For the treatment of the signs and symptoms ..."

< Back to Home

Register and Save Up to \$40 per Prescription* of MIRAPEX ER

You may be eligible to receive savings up to \$40 on each of your next 4 prescriptions of Once-daily MIRAPEX ER.

All you have to do is REGISTER TODAY!

*Up to 4 prescriptions.

This offer is not valid if you are enrolled in a Medicare, Medicaid, TriCare, or any other federal or state medical or pharmaceutical assistance program. This offer is not valid in Massachusetts or where prohibited by law.

The following questions are being asked by Boehringer Ingelheim Pharmaceuticals, Incorporated, so that we may advertise our products to you and provide you with information about them.

Please complete the following fields:

*Denotes required field.

First Name*

Last Name*

E-mail*

DOB* Month Year (You must be 18 years of age or older to enroll)

Address 1*

Address 2

City*

State* (Residents of MA are not eligible)

ZIP Code*

Are you a patient with Parkinson's disease or are you a caregiver?*

Patient Caregiver

How long ago were you or the person you care for diagnosed with Parkinson's disease?*

- Less than a year
- 1-3 years
- 3-5 years
- 5-7 years
- 7-9 years
- 10+ years

Are you or the person you care for currently taking medication for Parkinson's disease?*

Yes No

Which medication for Parkinson's disease are you taking (check all that apply)?*

- Mirapex ER® (pramipexole dihydrochloride) extended-release tablets
- Mirapex® (pramipexole dihydrochloride) tablets
- Requip® or ropinirole hydrochloride
- Requip®XL™ (ropinirole extended-release tablets)
- Stalevo® (carbidopa, levodopa, and entacapone)
- Azilect® (rasagiline)
- Sinemet® (carbidopa-levodopa) or any combination of carbidopa and levodopa
- Comtan® (entacapone)
- Eldepry® (selegiline hydrochloride) or Zelapar® (selegiline hydrochloride) Orally Disintegrating Tablets
- Symmetrel® (amantadine hydrochloride)
- Other

Are your prescriptions paid for in part or in full under any state or federally funded programs, including but not limited to Medicare, Medicaid, Medigap, VA, DoD, or TriCare?*

Yes No

Which kind of communication do you prefer?*

Postal mail E-mail

We respect your right to have personal and medical information kept confidential. Companies working with us will use the information you provide to send you information and help develop products, services, and programs. If we provide you with coupons or vouchers for our products, the information we receive from the pharmacy regarding their use will also be used for the same purposes.

We won't share your private information with anyone else—including mailing lists. Please check and sign below.

- By checking here, you indicate that you agree to the above and want us to use the information you have provided to send you information and offers relating to Parkinson's disease and MIRAPEX ER from time to time.
- By checking here, you indicate that you agree to the above and allow us to use your information to advertise Parkinson's disease-related and other products to you. Boehringer Ingelheim may also provide you with useful materials and contact you about health-related topics.

If you do not check either box, we will fill just this one-time request.

You are about to submit the above information. Please click "Submit" to verify that all answers are correct. If you need to make changes, click "Clear form" to do so.

[Clear form](#)

Trademarks are the property of their respective owners.



Indication:

MIRAPEX ER and MIRAPEX tablets are prescription medicines to treat the signs and symptoms of Parkinson's disease (PD).

Important Safety Information:

MIRAPEX ER or MIRAPEX may cause you to fall asleep without warning during daily activities, including talking, eating, and driving, which may result in accidents. Tell your doctor if you feel new or increased sleepiness while taking MIRAPEX ER or MIRAPEX. Do not drive a car, operate a machine, or do anything that needs you to be alert until you know how MIRAPEX ER or MIRAPEX affects you.

Before taking MIRAPEX ER or MIRAPEX, talk to your doctor about all of your medical conditions, including if you feel sleepy during the day. Tell your doctor if you drink alcohol or take other medications that make you drowsy, as these can increase the chance that MIRAPEX ER or MIRAPEX will make you feel sleepy or fall asleep when you should be awake.

When taking MIRAPEX ER or MIRAPEX, hallucinations (unreal visions, sounds or sensations) may occur and you may sometimes feel dizzy, nauseated, faint or sweaty when you sit up or stand quickly.

Some patients taking certain medicines to treat PD, including MIRAPEX ER and MIRAPEX, have reported problems, such as gambling, compulsive eating, compulsive buying, and increased sex drive. If you or your family members notice that you are developing unusual urges or behaviors, talk to your doctor.

The most common side effects in people taking MIRAPEX ER for early PD are sleepiness, nausea and vomiting, constipation, dizziness, tiredness, hallucinations, dry mouth, muscle spasms, and edema (swelling of the feet and ankles).

The most common side effects in people taking MIRAPEX ER who have later-stage PD are abnormal movements, nausea, constipation, hallucinations, headache, and anorexia.

The most common side effects in people taking MIRAPEX for PD are nausea, dizziness, sleepiness, constipation, hallucinations, insomnia, muscle weakness, confusion, and abnormal movements.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full Prescribing Information for [MIRAPEX ER](#) and [MIRAPEX](#).

This information is intended for U.S. residents only.

Included new privacy language: 1st sentence of privacy language ("The following ...") is here, the rest down below the questions.

Instead of saying "PD patient"

Instead of saying "PD medication"

Instead of saying "PD medication"

Changed order of these two questions about Medicare and Communication preference

New Privacy Language

New Fair Balance

[< Back to Home](#)

Register and Save Up to \$40 per Prescription* of MIRAPEX ER

You may be eligible to receive savings up to \$40 on each of your next 4 prescriptions of Once-daily MIRAPEX ER.

All you have to do is REGISTER TODAY!

*Up to 4 prescriptions.

This offer is not valid if you are enrolled in a Medicare, Medicaid, TriCare, or any other federal or state medical or pharmaceutical assistance program. This offer is not valid in Massachusetts or where prohibited by law.

The following questions are being asked by Boehringer Ingelheim Pharmaceuticals, Incorporated, so that we may advertise our products to you and provide you with information about them.

Please complete the following fields:

*Denotes required field.

First Name*

Last Name*

E-mail*

DOB* (You must be 18 years of age or older to enroll)

Address 1*

Address 2

City*

State* **(Residents of MA are not eligible)**

ZIP Code*

Are you a patient with Parkinson's disease or are you a caregiver?*

Patient Caregiver

How long ago were you or the person you care for diagnosed with Parkinson's disease?*

Less than a year
 1-3 years
 3-5 years
 5-7 years
 7-9 years
 10+ years

Are you or the person you care for currently taking medication for Parkinson's disease?*

Yes No

Which medication for Parkinson's disease are you taking (check all that apply)?*

- Mirapex ER® (pramipexole dihydrochloride) extended-release tablets
 Mirapex® (pramipexole dihydrochloride) tablets
 Requip® or ropinirole hydrochloride
 Requip®XL™ (ropinirole extended-release tablets)
 Stalevo® (carbidopa, levodopa, and entacapone)
 Azilect® (rasagiline)
 Sinemet® (carbidopa-levodopa) or any combination of carbidopa and levodopa
 Comtan® (entacapone)
 Eldepry® (selegiline hydrochloride) or Zelapar® (selegiline hydrochloride) Orally Disintegrating Tablets
 Symmetrel® (amantadine hydrochloride)
 Other

Are your prescriptions paid for in part or in full under any state or federally funded programs, including but not limited to Medicare, Medicaid, Medigap, VA, DoD, or TriCare?*

Yes No

Which kind of communication do you prefer?*

Postal mail E-mail

We respect your right to have personal and medical information kept confidential. Companies working with us will use the information you provide to send you information and help develop products, services, and programs. If we provide you with coupons or vouchers for our products, the information we receive from the pharmacy regarding their use will also be used for the same purposes.

We won't share your private information with anyone else—including mailing lists. Please check and sign below.

By checking here, you indicate that you agree to the above and want us to use the information you have provided to send you information and offers relating to Parkinson's disease and MIRAPEX ER from time to time.

By checking here, you indicate that you agree to the above and allow us to use your information to advertise Parkinson's disease-related and other products to you. Boehringer Ingelheim may also provide you with useful materials and contact you about health-related topics.

If you do not check either box, we will fill just this one-time request.

You are about to submit the above information. Please click "Submit" to verify that all answers are correct. If you need to make changes, click "Clear form" to do so.

[Clear form](#)

Trademarks are the property of their respective owners.



Indication:

MIRAPEX ER and MIRAPEX tablets are prescription medicines to treat the signs and symptoms of Parkinson's disease (PD).

Important Safety Information:

MIRAPEX ER or MIRAPEX may cause you to fall asleep without warning during daily activities, including talking, eating, and driving, which may result in accidents. Tell your doctor if you feel new or increased sleepiness while taking MIRAPEX ER or MIRAPEX. Do not drive a car, operate a machine, or do anything that needs you to be alert until you know how MIRAPEX ER or MIRAPEX affects you.

Before taking MIRAPEX ER or MIRAPEX, talk to your doctor about all of your medical conditions, including if you feel sleepy during the day. Tell your doctor if you drink alcohol or take other medications that make you drowsy, as these can increase the chance that MIRAPEX ER or MIRAPEX will make you feel sleepy or fall asleep when you should be awake.

When taking MIRAPEX ER or MIRAPEX, hallucinations (unreal visions, sounds or sensations) may occur and you may sometimes feel dizzy, nauseated, faint or sweaty when you sit up or stand quickly.

Some patients taking certain medicines to treat PD, including MIRAPEX ER and MIRAPEX, have reported problems, such as gambling, compulsive eating, compulsive buying, and increased sex drive. If you or your family members notice that you are developing unusual urges or behaviors, talk to your doctor.

The most common side effects in people taking MIRAPEX ER for early PD are sleepiness, nausea and vomiting, constipation, dizziness, tiredness, hallucinations, dry mouth, muscle spasms, and edema (swelling of the feet and ankles).

The most common side effects in people taking MIRAPEX ER who have later-stage PD are abnormal movements, nausea, constipation, hallucinations, headache, and anorexia.

The most common side effects in people taking MIRAPEX for PD are nausea, dizziness, sleepiness, constipation, hallucinations, insomnia, muscle weakness, confusion, and abnormal movements.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full Prescribing Information for [MIRAPEX ER](#) and [MIRAPEX](#).

This information is intended for U.S. residents only.

[< Back to Home](#)

Register and Save Up to \$40 per Prescription* of MIRAPEX ER

You may be eligible to receive savings up to \$40 on each of your next 4 prescriptions of Once-daily MIRAPEX ER.

All you have to do is REGISTER TODAY!

*Up to 4 prescriptions.

This offer is not valid if you are enrolled in a Medicare, Medicaid, TriCare, or any other federal or state medical or pharmaceutical assistance program. This offer is not valid in Massachusetts or where prohibited by law.

The following questions are being asked by Boehringer Ingelheim Pharmaceuticals, Incorporated, so that we may advertise our products to you and provide you with information about them.

Please complete the following fields:

*Denotes required field.

First Name*

Last Name*

E-mail*

DOB* (You must be 18 years of age or older to enroll)

Address 1*

Address 2

City*

State* **(Residents of MA are not eligible)**

ZIP Code*

Are you a patient with Parkinson's disease or are you a caregiver?*

Patient Caregiver

How long ago were you or the person you care for diagnosed with Parkinson's disease?*

Less than a year
 1-3 years
 3-5 years
 5-7 years
 7-9 years
 10+ years

Are you or the person you care for currently taking medication for Parkinson's disease?*

Yes No

Which medication for Parkinson's disease are you taking (check all that apply)?*

Mirapex ER® (pramipexole dihydrochloride) extended-release tablets
 Mirapex® (pramipexole dihydrochloride) tablets
 Requip® or ropinirole hydrochloride
 Requip® XL™ (ropinirole extended-release tablets)

Because you live in Massachusetts or because your prescriptions are paid for in part or in full under a state or federally funded program, including but not limited to Medicare, Medicaid, Medigap, VA, DoD, or TriCare, you are not eligible for the \$40 discount on prescriptions for MIRAPEX ER (pramipexole dihydrochloride) extended-release tablets.

changed placement of parentheses from after "extended-release" to after "dihydrochloride"

Yes No

Which kind of communication do you prefer?*

Postal mail E-mail

We respect your right to have personal and medical information kept confidential. Companies working with us will use the information you provide to send you information and help develop products, services, and programs. If we provide you with coupons or vouchers for our products, the information we receive from the pharmacy regarding their use will also be used for the same purposes.

We won't share your private information with anyone else—including mailing lists. Please check and sign below:

By checking here, you indicate that you agree to the above and want us to use the information you have provided to send you information and offers relating to Parkinson's disease and MIRAPEX ER from time to time.

By checking here, you indicate that you agree to the above and allow us to use your information to advertise Parkinson's disease-related and other products to you. Boehringer Ingelheim may also provide you with useful materials and contact you about health-related topics.

If you do not check either box, we will fill just this one-time request.

You are about to submit the above information. Please click "Submit" to verify that all answers are correct. If you need to make changes, click "Clear form" to do so.

[Clear form](#)

Trademarks are the property of their respective owners.



Indication:

MIRAPEX ER and MIRAPEX tablets are prescription medicines to treat the signs and symptoms of Parkinson's disease (PD).

Important Safety Information:

MIRAPEX ER or MIRAPEX may cause you to fall asleep without warning during daily activities, including talking, eating, and driving, which may result in accidents. Tell your doctor if you feel new or increased sleepiness while taking MIRAPEX ER or MIRAPEX. Do not drive a car, operate a machine, or do anything that needs you to be alert until you know how MIRAPEX ER or MIRAPEX affects you.

Before taking MIRAPEX ER or MIRAPEX, talk to your doctor about all of your medical conditions, including if you feel sleepy during the day. Tell your doctor if you drink alcohol or take other medications that make you drowsy, as these can increase the chance that MIRAPEX ER or MIRAPEX will make you feel sleepy or fall asleep when you should be awake.

When taking MIRAPEX ER or MIRAPEX, hallucinations (unreal visions, sounds or sensations) may occur and you may sometimes feel dizzy, nauseated, faint or sweaty when you sit up or stand quickly.

Some patients taking certain medicines to treat PD, including MIRAPEX ER and MIRAPEX, have reported problems, such as gambling, compulsive eating, compulsive buying, and increased sex drive. If you or your family members notice that you are developing unusual urges or behaviors, talk to your doctor.

The most common side effects in people taking MIRAPEX ER for early PD are sleepiness, nausea and vomiting, constipation, dizziness, tiredness, hallucinations, dry mouth, muscle spasms, and edema (swelling of the feet and ankles).

The most common side effects in people taking MIRAPEX ER who have later-stage PD are abnormal movements, nausea, constipation, hallucinations, headache, and anorexia.

The most common side effects in people taking MIRAPEX for PD are nausea, dizziness, sleepiness, constipation, hallucinations, insomnia, muscle weakness, confusion, and abnormal movements.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full Prescribing Information for [MIRAPEX ER](#) and [MIRAPEX](#).

This information is intended for U.S. residents only.



[Important Safety Information](#)

[Prescribing Information](#)

[< Back to Home](#)

Thank You for registering!

To get started, click on the button below to download your coupon. Then just print and take the coupon to your pharmacist when you fill your prescriptions for MIRAPEX ER. Please note that only one coupon may be redeemed per patient.

 **Print Your Coupon**
For up to \$40 in Savings on
MIRAPEX ER Now.



If you can't afford your MIRAPEX ER, our patient assistance program may be able to help. Call 1-800-556-8317

Indication:

MIRAPEX ER and MIRAPEX tablets are prescription medicines to treat the signs and symptoms of Parkinson's disease (PD).

Important Safety Information:

MIRAPEX ER or MIRAPEX may cause you to fall asleep without warning during daily activities, including talking, eating, and driving, which may result in accidents. Tell your doctor if you feel new or increased sleepiness while taking MIRAPEX ER or MIRAPEX. Do not drive a car, operate a machine, or do anything that needs you to be alert until you know how MIRAPEX ER or MIRAPEX affects you.

Before taking MIRAPEX ER or MIRAPEX, talk to your doctor about all of your medical conditions, including if you feel sleepy during the day. Tell your doctor if you drink alcohol or take other medications that make you drowsy, as these can increase the chance that MIRAPEX ER or MIRAPEX will make you feel sleepy or fall asleep when you should be awake.

When taking MIRAPEX ER or MIRAPEX, hallucinations (unreal visions, sounds or sensations) may occur and you may sometimes feel dizzy, nauseated, faint or sweaty when you sit up or stand quickly.

Some patients taking certain medicines to treat PD, including MIRAPEX ER and MIRAPEX, have reported problems, such as gambling, compulsive eating, compulsive buying, and increased sex drive. If you or your family members notice that you are developing unusual urges or behaviors, talk to your doctor.

The most common side effects in people taking MIRAPEX ER for early PD are sleepiness, nausea and vomiting, constipation, dizziness, tiredness, hallucinations, dry mouth, muscle spasms, and edema (swelling of the feet and ankles).

The most common side effects in people taking MIRAPEX ER who have later-stage PD are abnormal movements, nausea, constipation, hallucinations, headache, and anorexia.

The most common side effects in people taking MIRAPEX for PD are nausea, dizziness, sleepiness, constipation, hallucinations, insomnia, muscle weakness, confusion, and abnormal movements.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full Prescribing Information for [MIRAPEX ER](#) and [MIRAPEX](#).

This information is intended for U.S. residents only.

Introducing a TREATMENT OPTION for PARKINSON'S DISEASE



Ask your doctor or click here to learn more

Mirapex

pramipexole dihydrochloride tablets

EMAIL THIS PAGE PRINT THIS PAGE [MIRAPEX FOR ANOTHER CONDITION](#) [FOR HEALTHCARE PROFESSIONALS](#)

Type Size: [A](#) [A](#) [A](#)

- ▶ What Is Parkinson's Disease?
- ▶ Treatment of PD with MIRAPEX
- ▶ Talking to Your Doctor
- ▶ Tips for Living with PD
- ▶ Caring for Someone with PD
- ▶ Resources

FOR THE INITIAL AND LONG-TERM TREATMENT OF PARKINSON'S DISEASE (PD)

Plan to continue staying active

It's important to stay active, and MIRAPEX may help! In fact, studies show MIRAPEX may improve the ability to perform everyday activities, such as walking, dressing and handwriting. This means MIRAPEX may help you continue to enjoy your life... every day!

[Read More about MIRAPEX](#)



Registration button, registration page and all call-outs throughout the site leading to it will be removed.

Welcome to the Patient and Care Partner Resource for Parkinson's Disease

The diagnosis of Parkinson's disease (PD) may seem like a lot to handle at first. After all, people with Parkinson's disease sometimes have trouble with everyday activities.

MIRAPEX treats the signs and symptoms of idiopathic Parkinson's Disease. MIRAPEX may improve the ability to perform everyday activities and reduce resting tremor, a common symptom of Parkinson's disease.

Think about the things you like to do and want to continue to enjoy. Then talk to your doctor, because MIRAPEX and other medications may help.

MIRAPEX for Parkinson's Disease
Learn more about how MIRAPEX works.

[Learn more](#)

Talking to Your Doctor*
Find out how to help your doctor help you.

[Learn more](#)

Caring for Someone with Parkinson's Disease
Find out what you can do for a loved one.

[Learn more](#)

*Doctor portrayal

Please review new FB layout and the banner above only.



If you can't afford your MIRAPEX, our patient assistance program may be able to help. Call 1-800-556-8317

Important Information about MIRAPEX:

MIRAPEX is indicated for the treatment of the signs and symptoms of idiopathic Parkinson's disease (PD).

MIRAPEX may cause you to fall asleep without any warning, even while doing normal daily activities, such as driving. Before taking MIRAPEX, talk to your doctor if you drink alcohol or take other medications that make you drowsy as these can increase the chance that MIRAPEX will make you feel sleepy or fall asleep when you should be awake. When taking MIRAPEX, hallucinations (seeing, hearing, feeling, or tasting something that isn't there) may occur and you may sometimes feel dizzy, nauseated, faint or sweaty when you sit up or stand quickly.

In clinical trials for early PD, the most commonly reported side effects of MIRAPEX that were more frequent than with placebo are nausea, dizziness, sleepiness, difficulty falling asleep, weakness, and constipation. In clinical trials for advanced PD, the most commonly reported side effects of MIRAPEX that were more frequent than with placebo are low blood pressure when you sit up or stand quickly, involuntary movement, difficulty falling asleep, dizziness, accidental injury, hallucinations, and dream abnormalities. You should talk with your doctor if you experience these problems.

There have been reports of impulse control disorders and compulsive behaviors in patients taking certain medicines, including MIRAPEX. If you or your family members notice that you are experiencing new or increased gambling urges, increased sexual urges or other intense urges, such as compulsive shopping or eating, while taking MIRAPEX, talk to your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see [full Prescribing Information](#).

This information is intended for U.S. residents only.



Introducing a TREATMENT OPTION for PARKINSON'S DISEASE from Boehringer Ingelheim

[Learn More](#)



[EMAIL THIS PAGE](#) [PRINT THIS PAGE](#) [MIRAPEX FOR ANOTHER CONDITION](#) [FOR PATIENTS AND CARE PARTNERS](#)

Type Size: [A](#) [A](#) [A](#)

- [▶ MIRAPEX Benefits](#)
- [▶ Initial Monotherapy](#)
- [▶ Combination Therapy](#)
- [▶ Safety & Dosing](#)
- [▶ About Parkinson's Disease](#)
- [▶ Resources](#)

FOR THE INITIAL AND LONG-TERM TREATMENT OF PARKINSON'S DISEASE (PD)

IMPROVEMENT IN MOVEMENT

(Yet, Sam still has two left feet)

MIRAPEX improves motor symptoms of Parkinson's disease

[▶ Learn more](#)

MIRAPEX

- Significantly improves [motor symptoms](#) of Parkinson's disease¹
- Helps [patients maintain their everyday activities](#) at all stages of Parkinson's disease²
- Delays and reduces [motor complications](#) vs levodopa^{1,2}
- [Reserves the use of levodopa](#)^{1,2}
- Significantly reduces [levodopa-resistant tremor](#)³
- Provides dosing options for ease of [titration](#) to the following strengths (tid): 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 1.5 mg
- Has a well-established [safety and tolerability profile](#)³

MIRAPEX offers flexible dosing and titration

[▶ Learn more](#)

Learn about drug interaction considerations with MIRAPEX

[▶ Click here](#)

Print out Parkinson's disease treatment guidelines*

[▶ Click here](#)

* You will need Adobe® Reader® to download this file. If you don't have Adobe Reader, [click here](#) for a free and secure download.

Please review new FB layout and the banner above only.



Important Information about MIRAPEX:

MIRAPEX is indicated for the treatment of the signs and symptoms of idiopathic Parkinson's disease (PD).

Patients have reported falling asleep without perceived warning signs during activities of daily living, including operation of a motor vehicle, which sometimes resulted in accidents. Hallucinations and postural (orthostatic) hypotension may occur. In clinical trials for early PD, the most commonly reported side effects of MIRAPEX that were more frequent than with placebo are nausea (28% vs. 18%), dizziness (25% vs. 24%), somnolence (22% vs. 9%), insomnia (17% vs. 12%), asthenia (14% vs. 12%), and constipation (14% vs. 6%). In clinical trials for advanced PD, the most commonly reported side effects of MIRAPEX that were more frequent than with placebo are postural hypotension (53% vs. 48%), dyskinesia (47% vs. 31%), extrapyramidal syndrome (28% vs. 26%), insomnia (27% vs. 22%), dizziness (26% vs. 25%), accidental injury (17% vs. 15%), hallucinations (17% vs. 4%), and dream abnormalities (11% vs. 10%).

Patients and caregivers should be informed that impulse control disorders and compulsive behaviors have been reported in patients taking dopamine agonists, including MIRAPEX.

Please see [full Prescribing Information](#).

This information is intended for U.S. residents only.



[Privacy Statement](#) | [Terms of Use](#) | [Prescribing Information](#) | [References](#) | [Home](#) | [Site Map](#)